relativity of speed

borz vanni

August 10, 2026

Abstract

This document formalizes a mathematical extension of Special Relativity based on a macroscopic observer within a material medium. By replacing the vacuum cosmic constant c with a particle-and-medium-dependent function, we define the effective arrival velocity (

veffv sub eff end-sub eff). This model mathematically explains how neutrinos can arrive ahead of light in matter without violating causality, effectively redefining the local geometry perceived by the observer.

1 Introduction and Core Postulate

In the classical framework of relativity, the constant c represents the universal speed limit of spacetime in a vacuum. However, for a macroscopic observer measuring arrival times through a material medium, the concept of a limiting velocity must be redefined relative to the specific particle used as a probe.

We postulate that spacetime within matter exhibits an asymmetric geometry: light undergoes electromagnetic interactions that lengthen its path (scattering effects), whereas the neutrino (guided solely by the weak force or higher-dimensional physics) experiences a straight geodetic line (a geometric shortcut).

2 The Effective Velocity Formula

We define the **Effective Velocity of Arrival** (

veffv sub eff end-sub eff) for a generic particle x as:

veff(x)=ccos(x)nx1 lines; Line 1: v sub eff end-sub open paren x close paren equals the fraction with numerator c center dot cosine open paren theta sub x close paren and denominator n sub x end-fraction end-lines; eff()=cos()

Where the associated parameters are defined as:

- **c**: The universal limiting velocity of spacetime in a vacuum (300,000 km/s).
- nxn sub x

  nxn sub x : The refractive index or energetic slowing factor of the medium for particle x.
- (

) **cos**

  (

) **: The geometric shortcut factor, where**

  xtheta sub x represents the mean deviation angle relative to the macroscopic straight line.

2.1 Relative Parameter Assignment

For the macroscopic observer, the two primary particles behave according to the following mathematical properties:

$$\text{For the Neutrino ():} \quad n_\nu \approx 1, \quad \theta_\nu = 0^\circ \implies \cos(0^\circ) = 1 \implies v_{\text{eff}}(\nu) \approx c \tag{1}$$

$$\text{For Light (photon):} \quad n_{\text{light}} > 1, \quad \theta_{\text{light}} > 0^\circ \implies \cos(\theta_{\text{light}}) < 1 \implies v_{\text{eff}}(\text{light}) < c \tag{2}$$

2.2 Time-Advance Condition

The neutrino arrives ahead of light from the observer's perspective if and only if its effective velocity is strictly greater than that of the photon:

veff()¿veff(light)1n¿cos(light)nlight1 lines; Line 1: v sub eff end-sub open paren nu close paren is greater than v sub eff end-sub open paren light close paren the fraction with numerator 1 and denominator n sub nu end-fraction is greater than the fraction with numerator cosine open paren theta sub light end-sub close paren and denominator n sub light end-sub end-fraction end-lines; eff()¿eff(light)1¿cos(light)light

Since

1 , the condition reduces to the inequality which is always verified within matter:

1¿cos(light)nlight1 lines; Line 1: 1 is greater than the fraction with numerator cosine open paren theta sub light end-sub close paren and denominator n sub light end-sub end-fraction end-lines; 1¿cos(light)light

3 Rewriting Einstein's Formulae

We introduce the **Effective Lorentz Factor** (

effgamma sub eff end-sub eff) for a body or particle moving at a local velocity v within the medium, embedding the interchangeability of the limiting constant:

eff(x)=11v2veff(x)21 lines; Line 1: gamma sub eff end-sub open paren x close paren equals the fraction with numerator 1 and denominator the square root of 1 minus the fraction with numerator v squared and denominator v sub eff end-sub open paren x close paren squared end-fraction end-root end-fraction end-lines; eff()=112eff()2

We now apply this substitution to the fundamental equations of Relativistic Energy and Mechanics.

3.1 Effective Relativistic Energy

The total energy calculated by the observer for the particle within the medium becomes dependent on its specific geometric arrival limit:

$$E = \gamma_{\text{eff}}(x) \cdot m \cdot v_{\text{eff}}(x)^2 = \frac{m \cdot v_{\text{eff}}(x)^2}{\sqrt{1 - \frac{v^2}{v_{\text{eff}}(x)^2}}} \tag{3}$$

3.2 Relativistic Momentum

The mechanical momentum or impulse of the particle relative to the observer transforms into:

$$p = \gamma_{\text{eff}}(x) \cdot m \cdot v = \frac{m \cdot v}{\sqrt{1 - \frac{v^2}{v_{\text{eff}}(x)^2}}} \tag{4}$$

This rewriting allows the neutrino to move at velocities greater than local light (

¿eff

(

light
)) while maintaining a real and calculable momentum, since the denominator utilizes eff
(
)
.

3.3 Relative Time Dilation

The proper time interval t' measured by the moving particle relative to the macroscopic observer's time t becomes:

t=teff(x)=t1v2veff(x)21 lines; Line 1: delta t prime equals the fraction with numerator delta t and denominator gamma sub eff end-sub open paren x close paren end-fraction equals delta t center dot the square root of 1 minus the fraction with numerator v squared and denominator v sub eff end-sub open paren x close paren squared end-fraction end-root end-lines; =eff()=12eff()2

4 Conclusions

This model demonstrates that the constant c is not violated, but is rather rewritten as a geometric

. The interchangeability between light and neutrinos in the formulae depends purely on the observer's state of necessity and the chemical-physical interaction type of the crossed material, thus resolving macroscopic arrival-time paradoxes.